

A Liouville Type Theorem for Steady-State Navier-Stokes Equations

G Seregin

University of Oxford, UK and PDMI, RAS, Russia

November 8, 2016

Abstract

A Liouville type theorem is proven for the steady-state Navier-Stokes equations. It follows from the corresponding theorem on the Stokes equations with the drift. The drift is supposed to belong to a certain Morrey space.

1 The Main Result

The classical Liouville type theorem for the stationary Navier-Stokes equations can be stated as follows: show that any bounded solution to the system

$$u \cdot \nabla u - \Delta u = \nabla p, \quad \operatorname{div} u = 0 \quad (1.1)$$

is constant. This problem has not solved yet and even it is not clear if it has a positive answer.

Another popular problem is to show that any solution to system (1.1), satisfying two conditions:

$$\int_{\mathbb{R}^3} |\nabla u|^2 dx < \infty. \quad (1.2)$$

and

$$u(x) \rightarrow 0 \quad \text{as} \quad |x| \rightarrow \infty, \quad (1.3)$$

is identically equal to zero. Unfortunately, it is still unknown whether the this statement is true or not.

However, some attempts have been made to solve above or related problems. One of the best results in that direction can be found in [4] where it is shown that the assumption

$$u \in L_{\frac{9}{2}}(\mathbb{R}^3) \quad (1.4)$$

implies $u = 0$. Very recently, condition (1.4) has been improved logarithmically in [3].

Another set of admissible functions for solutions to (1.1), in which the Liouville type theorem is valid, has been described in [9]. To be precise, any solution to (1.1), obeying the inclusion

$$u \in L_6(\mathbb{R}^3) \cap BMO^{-1}(\mathbb{R}^3), \quad (1.5)$$

is identically equal to zero.

For more Liouville type results, we refer the reader to interesting papers [6], [7], [2], and [1] and references there.

Our short note is inspired by paper [8] by Nazarov-Uraltseva about properties of solutions to elliptic and parabolic linear equations with divergence free drift. Although their approach works for scalar equations only, similar assumptions on the drift occur in the vectorial case as well. We formulate our result as a statement of the linear theory, considering the following steady-state Stokes system with the drift

$$u \cdot \nabla v - \Delta v = \nabla q, \quad \operatorname{div} v = 0, \quad \operatorname{div} u = 0. \quad (1.6)$$

Theorem 1.1. *Suppose that smooth functions u and v satisfy (1.6) and two additional conditions:*

$$M := \sup_{R>0} R^{1-\frac{3}{q}} \|u\|_{L^{q,\infty}(B(R))} < \infty \quad (1.7)$$

with $3/2 < q \leq 3$ and

$$N := \sup_{R>0} R^{\frac{1}{2}-\frac{3}{s}} \|v\|_{s,B(R)} < \infty \quad (1.8)$$

with $2 \leq s \leq 6$. Then $v \equiv 0$ in $\mathbb{R}^3$.

Here, $L^{q,\infty}(\Omega)$ stands for a weak Lebesgue space, which is a particular Lorentz space $L^{q,r}(\Omega)$ and $L^{q,q}(\Omega) = L_q(\Omega)$ is a usual Lebesgue space.

It is an interesting question to understand difference between above conditions (1.4) and (1.7), (1.8) for $u = v$. To this end, assume that there exists a divergence free field u having the following bound from above

$$|u(x)| \leq \frac{1}{|x'| + 1} \frac{1}{(|x_3| + 1)^{\frac{2}{9}}}.$$

Then condition (1.7) holds if q is not equal to 2 and condition (1.8) holds with $s = 6$ while condition (1.4) is violated.

2 Proof of Main Result

2.1 Caccioppoli Type Inequality

Let $0 < R < 2$ and let a non-negative cut-off function $\varphi \in C_0^\infty(B(R))$ satisfy the following properties: $\varphi(x) = 1$ in $B(r)$, $\varphi(x) = 0$ out of $B(R)$, and $|\nabla\varphi(x)| \leq c/(R-r)$ for any $1 \leq r < R \leq 2$. We let $\bar{u} = v - [v]_{B(R)}$, where $[v]_{B(R)}$ is the mean value of v over the ball of radius R centred at the origin.

A given exponent q , satisfying conditions of Theorem 1.1, see (1.7), one can find a constant $c_0(q)$ and a function w_R that is smooth in $B(2)$, vanishes outside $B(R)$ and satisfies the identity $\operatorname{div} w_R = \nabla\varphi \cdot \bar{v}$ and the inequality

$$\|\nabla w_R\|_{L^{2q',2}(B(R))} \leq c_0 \|\nabla\varphi \cdot \bar{v}\|_{L^{2q',2}(B(R))} \leq \frac{c_0}{R-r} \|\bar{v}\|_{L^{2q',2}(B(R))}. \quad (2.1)$$

Moreover, by interpolation and Hardy-Littlewood-Sobolev inequality, we also have a bound for the right hand side of (2.1):

$$\|\bar{v}\|_{L^{2q',2}(B(R))} < c(q) \|\bar{v}\|_{2,B(R)}^{1-\frac{3}{2q}} \|\nabla v\|_{2,B(R)}^{\frac{3}{2q}}. \quad (2.2)$$

Now, let us test the first equation in (1.6) with the function $\varphi\bar{v} - w_R$, integrate by parts in $B(R)$, and find the following identity

$$\begin{aligned} \int_{B(R)} \varphi |\nabla v|^2 dx &= - \int_{B(R)} \nabla v : (\nabla\varphi \otimes \bar{v}) dx + \int_{B(R)} \nabla w_R : \nabla v dx + \\ &- \int_{B(R)} (u \cdot \nabla v) \cdot \varphi \bar{v} dx + \int_{B(R)} (u \cdot \nabla v) \cdot w_R dx = I_1 + I_2 + I_3 + I_4. \end{aligned}$$

I_1 can be estimated easily. As a result, the below bound is valid:

$$|I_1| \leq \frac{c}{R-r} \|\nabla v\|_{2,B(R)} \|\bar{v}\|_{2,B(R)}.$$

As to I_2 , by Hölder inequality, we have

$$\begin{aligned} |I_2| &\leq \|\nabla v\|_{2,B(R)} \|\nabla w_R\|_{2,B(R)} = \|\nabla v\|_{2,B(R)} \|\nabla w_R\|_{L^{2,2}(B(R))} \leq \\ &\leq \|\nabla v\|_{2,B(R)} \|\nabla w_R\|_{L^{2q',2}(B(R))} \|1\|_{L^{2q,\infty}(B(R))} \leq \\ &\leq cR^{\frac{3}{2q}} \|\nabla v\|_{2,B(R)} \|\nabla w_R\|_{L^{2q',2}(B(R))}. \end{aligned}$$

Now, taking into account (2.1) and (2.2), one can derive from the latter estimate the following:

$$\begin{aligned} |I_2| &\leq c \|\nabla v\|_{2,B(R)} \frac{R^{\frac{3}{2q}}}{R-r} \|\bar{v}\|_{2,B(R)}^{1-\frac{3}{2q}} \|\nabla v\|_{2,B(R)}^{\frac{3}{2q}} \leq \\ &\leq c \frac{R}{R-r} \|\nabla v\|_{2,B(R)}^{1+\frac{3}{2q}} \left(\frac{1}{R} \|\bar{v}\|_{2,B(R)} \right)^{1-\frac{3}{2q}}. \end{aligned}$$

Let us start evaluation of I_3 with integration by parts that gives

$$I_3 = \frac{1}{2} \int_{B(R)} |\bar{v}|^2 u \cdot \nabla \varphi dx.$$

Hence,

$$\begin{aligned} |I_3| &\leq \frac{c}{R-r} \|u\|_{L^{q,\infty}(B(R))} \|\bar{v}\|^2_{L^{q',1}(B(R))} \leq \\ &\leq \frac{c}{R-r} \|u\|_{L^{q,\infty}(B(R))} \|\bar{v}\|^2_{L^{2q',2}(B(R))} \leq \\ &\leq \frac{c}{R-r} \|u\|_{L^{q,\infty}(B(R))} \|\bar{v}\|_{2,B(R)}^{2(1-\frac{3}{2q})} \|\nabla v\|_{2,B(R)}^{2\frac{3}{2q}} \leq \\ &\leq c \frac{R}{R-r} M_0 \left(\frac{1}{R^2} \|\bar{v}\|_{2,B(R)}^2 \right)^{1-\frac{3}{2q}} \|\nabla v\|_{2,B(R)}^{2\frac{3}{2q}}, \end{aligned}$$

where

$$M_0 = \sup_{0 < R < 2} R^{1-\frac{3}{2q}} \|u\|_{L^{q,2}(B(R))}.$$

The last term can be estimated in a similar way. Indeed, integrating by parts and applying Hölder inequality,

$$\begin{aligned}
|I_4| &= \left| \int_{B(R)} (u \cdot \nabla w_R) \cdot \bar{v} dx \right| \leq \|u\|_{L^q, \infty(B(R))} \|\nabla w_R\| \|\bar{v}\|_{L^{q', 1}(B(R))} \leq \\
&\leq \|u\|_{L^q, \infty(B(R))} \|\nabla w_R\|_{L^{2q', 2}(B(R))} \|\bar{v}\|_{L^{2q', 2}(B(R))} \leq \\
&\leq \frac{c}{R-r} \|u\|_{L^q, \infty(B(R))} \|\bar{v}\|^2_{L^{q', 1}(B(R))}.
\end{aligned}$$

The right hand side of the latter inequality has been already estimated. Hence, we find

$$|I_4| \leq c \frac{R}{R-r} M_0 \left(\frac{1}{R^2} \|\bar{v}\|_{2, B(R)}^2 \right)^{1 - \frac{3}{2q}} \|\nabla v\|_{2, B(R)}^{2 \frac{3}{2q}}.$$

Summarising four above estimates, we show

$$\begin{aligned}
f(r) &\leq c \frac{R}{R-r} f^{\frac{1}{2}}(R) \left(\frac{1}{R^2} \|\bar{v}\|_{2, B(R)}^2 \right)^{\frac{1}{2}} + \\
&+ c \frac{R}{R-r} (f(R))^{\frac{1}{2}(1 + \frac{3}{2q})} \left(\frac{1}{R^2} \|\bar{v}\|_{2, B(R)}^2 \right)^{\frac{1}{2}(1 - \frac{3}{2q})} + \\
&+ c \frac{R}{R-r} M_0 \left(\frac{1}{R^2} \|\bar{v}\|_{2, B(R)}^2 \right)^{1 - \frac{3}{2q}} (f(R))^{\frac{3}{2q}},
\end{aligned}$$

where

$$f(R) = \|\nabla v\|_{2, B(R)}^2.$$

For any $1 \leq R \leq 2$,

$$\frac{1}{R^2} \|\bar{v}\|_{2, B(R)}^2 \leq \|\widehat{v}\|_{2, B(2)}^2$$

with $\widehat{v} = v - [v]_{B(2)}$.

Given $\varepsilon > 0$, applying Young inequality, we find

$$f(r) \leq \varepsilon f(R) + c(M_0, q, \varepsilon) \|\widehat{v}\|_{2, B(2)}^2 \left(\frac{1}{(R-r)^2} + \frac{1}{(R-r)^{\kappa_1}} + \frac{1}{(R-r)^{\kappa_2}} \right)$$

for any $1 \leq R \leq 2$, where

$$\kappa_1 = \frac{1}{\frac{1}{2}(1 - \frac{3}{2q})}, \quad \kappa_2 = \frac{1}{1 - \frac{3}{2q}}.$$

As it has been shown in [5], there exists a positive number ε depending on M_0 and q only such that

$$\int_{B(1)} |\nabla v|^2 dx \leq c(M_0, q) \int_{B(2)} |v - [v]_{B(2)}|^2 dx.$$

It is known that the Navier-Stokes equations are invariant with respect to the shift and the scaling of the form

$$v(x, t) \rightarrow \lambda v(\lambda x, \lambda^2 t), \quad q(x, t) \rightarrow \lambda^2 q(\lambda x, \lambda^2 t).$$

This allows us to get the required Caccioppoli type inequality

$$\int_{B(x_0, R)} |\nabla v|^2 dx < c(M, q) \frac{1}{R^2} \int_{B(x_0, 2R)} |v - [v]_{B(x_0, 2R)}|^2 dx \quad (2.3)$$

being valid for any $R > 0$ and $x_0 \in \mathbb{R}^3$.

2.2 Proof of Theorem 1.1

We can put $x_0 = 0$ and use the following simple inequality

$$\frac{1}{R^2} \int_{B(2R)} |v - [v]_{B(2R)}|^2 dx \leq c \frac{1}{R^2} \int_{B(2R)} |v|^2 dx \leq \frac{1}{R^{2(\frac{3}{s} - \frac{1}{2})}} \|v\|_{B(2R)}^2 \leq c N^2$$

for any $R > 0$. Passing $R \rightarrow \infty$, we conclude that

$$\int_{\mathbb{R}^3} |\nabla v|^2 dx < \infty$$

The rest of the proof is the same as in [9].

References

- [1] Chae, D., Liouville-Type Theorem for the Forced Euler Equations and the Navier-Stokes Equations. Commun. Math. Phys. 326: 37-48 (2014).

- [2] Chae, D., Yoneda, T., On the Liouville theorem for the stationary Navier-Stokes equations in a critical space, *J. Math. Anal. Appl.* 405 (2013), no. 2, 706-710.
- [3] Chae, G., Wolf, J., On Liouville type theorems for the steady Navier-Stokes equations in R^3 , arXiv:1604.07643.
- [4] Galdi, G. P. An introduction to the mathematical theory of the Navier-Stokes equations. Steady-state problems. Second edition. Springer Monographs in Mathematics. Springer, New York, 2011. xiv+1018 pp.
- [5] Giaquinta, M., Multiple integrals in the calculus of variations and non-linear elliptic systems. *Annals of Mathematics Studies*, 105. Princeton University Press, Princeton, NJ, 1983. vii+297 pp.
- [6] Gilbarg, D., Weinberger, H. F. Asymptotic properties of steady plane solutions of the Navier-Stokes equations with bounded Dirichlet integral, *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*(4) 5 (1978), no. 2, 381-404.
- [7] Koch, G., Nadirashvili, N., Seregin, G., Sverak, V., Liouville theorems for the Navier-Stokes equations and applications, *Acta Mathematica*, 203 (2009), 83–105.
- [8] Nazarov, A. I. and Uraltseva, N. N., The Harnack inequality and related properties for solutions to elliptic and parabolic equations with divergence-free lower-order coefficients, *St. Petersburg Mathematical Journal*, 2012, 23:1, 93–115.
- [9] Seregin, G., Liouville type theorem for stationary Navier-Stokes equations, *Nonlinearity*, 29 (2016), 2191–2195.
- [10] Stein, Elias M. Singular integrals and differentiability properties of functions. *Princeton Mathematical Series*, No. 30 Princeton University Press, Princeton, N.J. 1970 xiv+290 pp.